

JB Dental Pediatrics Information



Pediatric Treatment Notice

We know that parents are most concerned about their children's oral health. It is our common responsibility with you to protect children's oral health and cultivate correct oral hygiene habits. During treatment, your child may react nervously and resist due to the new environment. Thus, it is necessary for the company and positive encouragement of the parents to cooperate with our dentist, so that the child can undergo dental treatment safely smoothly. We have listed recommendations below, which we request that you and your children understand and accept before visiting children's dentistry.

Pre-treatment Precautions and Recommendations

1. Before accompanying your child to the dental clinic, it is advised that the child wears loose clothes and bring a set of clean clothes to avoid discomfort during the treatment process. If the child's clothes are splashed or dirty during the treatment, the child can immediately change.
2. Before seeing a doctor, please ensure that our child has brushed their teeth well to prevent food residues after meals from affecting the doctor's inspection and the effect of fluoride application. At your request, the doctor can also check whether the child's teeth brushing needs to be strengthened.
3. It is recommended that the child has not eaten for two hours for sleep therapy to avoid the child vomiting due to nervousness, crying, and oral sensitivity.
4. If your child has a history of drug allergies or diseases, please inform our medical team in advance.
5. Parents can help children build a trusting and friendly relationship with dentists or dental clinics and reduce children's resistance to unfamiliar doctors and clinic environments by being present and comforting the child.
6. It is recommended to encourage children to see a doctor with encouragement and rewards instead of scolding, and avoid words such as tooth extraction, injection, pain, and tooth drilling that deepen children's fear of nervousness.
7. Across from the front desk, JB Dental provides a fun and safe waiting area for children to help them ease their mood when seeing a doctor.

Our Approach to Pediatric Dentistry

1. At the beginning of the treatment, the doctor will establish a friendly relationship with the child, and kindly explain the tools used in the diagnosis and treatment to the child, so as to alleviate the child's fear of unfamiliar treatment tools and procedures.
2. We may adopt behavioral guidance depending on the. If the child does not cooperate with the diagnosis and treatment behavior, the doctor may raise their voice to ensure that the child cooperates with the doctor. This may be necessary for the child to understand the correct and safe behavior. Parents can rest assured that the doctors can handle children.
3. If the doctor asks the parents to leave and let them conduct the treatment independently, parents will be requested to cooperate with the doctor's instructions to facilitate the treatment.
4. Younger children are prone to crying and it may be difficult for them to cooperate with the doctor's instructions. At this time, the doctor may put their hands on their mouths to calm the children down and ensure that they pay attention to the doctor's instructions.
5. For children who refuse to cooperate with the treatment, the doctor will seek the consent of the parents to physically restrain the child's body movement using a restrain board to prevent the child from being scratched by sharp diagnostic tools due to movement. If the parents are worried about the restraint board/restraint, they can discuss with the doctor to evaluate whether to use sleep therapy.

Post-treatment Recommendations

1. After the visit, parents are requested to follow the doctor's instructions, and do not let the child eat immediately after.
2. If anesthesia is administered during the treatment, within two hours after the diagnosis and treatment, please remind the child to avoid repeatedly licking and biting the anesthetized part to avoid pain after the anesthesia dissipates.
3. (If the child is hungry and wants to eat, but the anesthetic has not returned, he can drink juice first or eat food that does not need to be chewed, such as pudding)
4. Please use positive and encouraging ways to increase children's confidence and courage to see a doctor in the future.
5. Parents are requested to take their children to the dental clinic for return visits on a regular basis, so as to avoid delaying the diagnosis and treatment and affecting the treatment results.
6. If you have any questions, please feel free to contact us.

